


Sec 13 HW - Implicit Differentiation

1. $x^2 + y^2 = 1$ is not a function, but it can be written as two fens:

$$y^2 = 1 - x^2 \rightarrow y_1 = \sqrt{1-x^2}, \quad y_2 = -\sqrt{1-x^2}$$


$$\leftarrow y_1 = \sqrt{1-x^2} \rightarrow y_1' = \frac{1}{2}(1-x^2)^{-1/2}(-2x)$$

$$\leftarrow y_2 = -\sqrt{1-x^2}$$

$$y_1' = \frac{-x}{\sqrt{1-x^2}}$$

$$y_2' = \frac{+x}{\sqrt{1-x^2}}$$

Are these the same as what implicit differentiation give?

$$x^2 + y^2 = 1$$

$$2x + 2y \frac{dy}{dx} = 0 \rightarrow \frac{dy}{dx} = \frac{-2x}{2y} =$$

$$\boxed{\frac{dy}{dx} = \frac{-x}{y}} = \frac{-x}{\pm\sqrt{1-x^2}} = \frac{\mp x}{\sqrt{1-x^2}} = \begin{matrix} y_1' \\ y_2' \end{matrix}$$

2a) $x^2 + 3y^2 = 6$

$$2x + 6y \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} = \frac{-2x}{6y} = \frac{-x}{3y}$$

$$\boxed{\frac{dy}{dx} = \frac{-x}{3y}}$$

b) $9x - x^2y^2 = 2xy$

Requires product rule:

$$9 - 2xy^2 - x^2 \cdot 2y \frac{dy}{dx} = 2y + 2x \frac{dy}{dx}$$



$$2b) \quad 9 - 2xy^2 - 2y = 2x^2y \frac{dy}{dx} + 2x \frac{dy}{dx}$$

$$9 - 2xy^2 - 2y = (2x^2y + 2x) \frac{dy}{dx}$$

$$\left| \frac{dy}{dx} = \frac{9 - 2xy^2 - 2y}{2x^2y + 2x} \right|$$

$$c) \quad 3xy - \frac{y}{3} = 2x^{-1}$$

$$3y + 3x \frac{dy}{dx} - \frac{1}{3} \frac{dy}{dx} = -2x^{-2}$$

$$\left(3x - \frac{1}{3} \right) \frac{dy}{dx} = -2x^{-2} - 3y$$

$$\frac{dy}{dx} = \frac{-2x^{-2} - 3y}{3x - \frac{1}{3}} = \frac{\frac{-2}{x^2} - 3y}{3x - \frac{1}{3}} \quad \text{LCD} = 3x^2$$

$$\frac{dy}{dx} = \frac{-3x^2}{-3x^2} \cdot \frac{\frac{-2}{x^2} - 3y}{3x - \frac{1}{3}} = \frac{6 + 9x^2y}{-9x^3 + x^2} \quad (\text{book has negative factored out})$$

$$f. \quad 3x^2 - 4y^3 + 3 = \sqrt{5xy}$$

$$6x - 12y^2 \frac{dy}{dx} = \frac{1}{2} (5xy)^{-1/2} \cdot \left(5 + \frac{dy}{dx} \right)$$

$$6x - 12y^2 \frac{dy}{dx} = \frac{5}{2} \left(\frac{1}{\sqrt{5xy}} \right) + \frac{1}{\sqrt{5xy}} \frac{dy}{dx}$$

$$6x - \frac{5}{2\sqrt{5xy}} = \left(12y^2 + \frac{1}{\sqrt{5xy}} \right) \frac{dy}{dx}$$


Sec 13 con'd

#2f

$$\frac{dy}{dx} = \frac{6x - \frac{5}{2\sqrt{5xy}}}{12y^2 + \frac{1}{\sqrt{5xy}}} \quad (\text{same as book, bit different form})$$

#3.

$$y^2 - x^2 = 16 \quad \text{hyperbola}$$



Eqn. of tangent line at $(2, 2\sqrt{5})$ is found:

$$2y \frac{dy}{dx} - 2x = 0$$

$$\left. \frac{dy}{dx} = \frac{x}{y} \right|_{(2, 2\sqrt{5})} = \frac{2}{2\sqrt{5}} = \frac{1}{\sqrt{5}} = \text{slope}$$

$$y - 2\sqrt{5} = \frac{1}{\sqrt{5}}(x - 2)$$

6.

$$e^{xy} = x$$

$$\left. \frac{dy}{dx} \right|_{x=3}$$

Find slope at $x=3$, that is,

(you'll need y value, so plug $x=3$ into $e^{xy} = x$ first)

$$e^{3y} = 3 \rightarrow \ln e^{3y} = \ln 3 \rightarrow 3y = \ln 3$$

$$\rightarrow y = \frac{\ln 3}{3} \text{ when } x=3$$

Now

nd $\frac{dy}{dx}$:

$$e^{xy} \left(y + x \frac{dy}{dx} \right) = 1 \rightarrow e^{xy} y + x e^{xy} \frac{dy}{dx} = 1$$

$$xe^{xy} \frac{dy}{dx} = 1 - ye^{xy} \rightarrow \boxed{\frac{dy}{dx} = \frac{1 - ye^{xy}}{xe^{xy}}}$$

$$\text{So } \left. \frac{dy}{dx} \right|_{(3, \ln 3)} = \frac{1 - \ln 3 e^{3 \ln 3}}{3 e^{3 \ln 3}}$$

book is wrong

$$= \frac{1 - \ln 3 \cdot e^{\ln 9}}{3 e^{\ln 9}} = \frac{1 - \ln 3 \cdot 9}{3 \cdot 9} = \boxed{\frac{1 - 9 \ln 3}{27}}$$

$$10. \quad y^2 = x \rightarrow 2y \frac{dy}{dx} = 1 \rightarrow \boxed{\frac{dy}{dx} = \frac{1}{2y}}$$

$$a) \text{ So } \frac{d^2 y}{dx^2} = -1(2y)^{-2} \cdot 2 \frac{dy}{dx} \leftarrow \begin{array}{l} \text{Chain rule} \\ \downarrow \\ \text{implicit} \end{array}$$

$$\text{thus } \frac{d^2 y}{dx^2} = \frac{-2}{2^2 y^2} \frac{dy}{dx} \leftarrow \text{Now substitute}$$

$$\frac{d^2 y}{dx^2} = \frac{-1}{2y^2} \cdot \frac{1}{2y} = \boxed{\frac{-1}{4y^3}}$$

b) Find $d^3 y / dx^3$ by quotient or power rule:

$$\text{Quotient rule: } \frac{d^3 y}{dx^3} = \frac{0(4y^3) - (-1)(12y^2 dy/dx)}{(4y^3)^2} = \frac{12y^2 dy/dx}{16y^6}$$

$$= \frac{3 dy/dx}{4y^4} \xrightarrow{\text{substitute } dy/dx} = \frac{3 \left(\frac{1}{2y} \right)}{4y^4} = \boxed{\frac{3}{8y^5}}$$

Power rule

$$\frac{d^2 y}{dx^2} = -(4y^3)^{-1} \rightarrow \frac{d^3 y}{dx^3} = +(4y^3)^{-2} \left(12y^2 \frac{dy}{dx} \right)$$

$$= \frac{12y^2}{(4y^3)^2} \frac{dy}{dx} = \frac{12y^2}{16y^6} \frac{dy}{dx} = \frac{3}{4y^4} \frac{dy}{dx}$$

Substituting $\frac{dy}{dx}$

$$\frac{d^3 y}{dx^3} = \frac{3}{4y^4} \cdot \frac{1}{2y} = \boxed{\frac{3}{8y^5}}$$

#8

For a certain product we have the following functions implicit in q , the first because it is not explicitly solved for C , the second, not explicitly solved for R .

Cost
Revenue

$$C^2 = q^2 + 100\sqrt{q} + 100$$

$$900(q-4)^2 + R^2 = 25,500$$

Marg. Cost
 $C'(q)$

$$2C \frac{dC}{dq} = 2q + \frac{1}{2} \cdot 100 q^{-1/2} + 0$$

$$2C \frac{dC}{dq} = 2q + \frac{50}{\sqrt{q}} \rightarrow \frac{dC}{dq} = \frac{2q}{2C} + \frac{50}{2C\sqrt{q}} = \boxed{\frac{q}{C} + \frac{25}{C\sqrt{q}}}$$

Marginal cost $C'(q)$

Marg. Rev
 $R'(q)$

$$1800(q-4) + 2R \frac{dR}{dq} = 0$$

$$\frac{dR}{dq} = \frac{-1800(q-4)}{2R} = \boxed{\frac{-900(q-4)}{R}}$$

marginal revenue function $R'(q)$.

To find $C'(5) + R'(5)$ we need to find what $C + R$ would be at $q = 5$, since the marginal functions are not explicit.

① Cost at $q = 5$: $C^2 = 5^2 + 100\sqrt{5} + 100$

$$C = \sqrt{25 + 100\sqrt{5}} = \sqrt{25(5 + 4\sqrt{5})} = 5\sqrt{5 + 4\sqrt{5}}$$

$$C'(q) = \frac{q\sqrt{q} + 25}{C\sqrt{q}} = \frac{5\sqrt{5} + 25}{5\sqrt{5+4\sqrt{5}}\sqrt{5}} = \frac{\sqrt{5} + 5}{\sqrt{5+4\sqrt{5}}\sqrt{5}}$$

Multiply top + bottom by $\sqrt{5}$:

$$C'(q) = \frac{\sqrt{5} + 5 \cdot \sqrt{5}}{\sqrt{5+4\sqrt{5}}\sqrt{5} \cdot \sqrt{5}} = \frac{5 + 5\sqrt{5}}{\sqrt{5+4\sqrt{5}} \cdot 5} = \frac{1 + \sqrt{5}}{\sqrt{5+4\sqrt{5}}}$$

A calculator tells $\approx .87$, or a rate of .87 / 100 items

② Revenue at $q = 5$: $900(5-4)^2 + R^2 = 25,500$
 $R^2 = 25,500 - 900 = 24,600$
 $R = \sqrt{24,600} = 10\sqrt{246}$

Then $R'(5) = \frac{-900(5-4)}{10\sqrt{246}} = \frac{-90}{\sqrt{246}}$ or -5.74 rate / 100 items

Meaning: When production is 500 units, costs are increasing at a rate of 87¢/next 100 items

When 500 units are sold, revenue for next 100 decreases by \$5.74