### **Summary of Critical Values, Derivative Tests and Concavity**

If f'(c) = 0, or if f'(c) does not exist (DNE), then c is a critical number. The function could have a local extreme at c (including a nondifferentiable cusp or corner) or it could have an *inflection point*.

Before invoking the first derivative test to see if the derivative changes sign, we could go straight to the second derivative test, testing f''(c) and drawing conclusions as follows:

- If f''(c) > 0, the function is concave up at c because the trend of the slope of the tangent is to increase. f(c) is a local min.
- If f''(c) < 0, the function is concave down at c because the trend of the slope of the tangent is to decrease. f(c) is a local max.
- If f''(c) = 0, we could have a local max or a local min, **or** we could have an inflection point. How do we decide what it is? There are a couple of things we can do:

### **EITHER**

1. Resort to the first derivative test, checking values on either side of c to see if f'(c) changes sign.

If it does, we have a local extreme at x = c.

If it doesn't, we have an inflection point at x = c.

### OR

2. Stay with the *second derivative test*, testing for a value on either side of c to see if f''(x) changes sign.

If f''(x) < 0 to the left of c and > 0 to the right of c (or if it is > 0 to the left and < 0 to the right), then c is clearly an inflection point because concavity has changed.

If f''(x) is > 0 on both sides then x = c is a local min

If f''(x) < 0 on both sides, then x = c is a local max.

Following are two examples that are illustrative because the functions are so simple you can already sketch them. Observe the derivative tests vis-a-vis the sketches to lock in your understanding.

# Example 1: Sketch the function $f(x) = x^4$

 $f'(x) = 4x^3 = 0$  at x = 0;  $f''(x) = 12x^2 = 0$  at x = 0, also. What kind of critical point, then, is c = 0? There are two ways to find out:

First derivative test: Checking values into f'(x) on either side of 0,  $f'(-1) = 4(-1)^3 = -4$  and  $f'(1) = 4(1)^3 = 4$ . Because f' changes sign, negative to positive, c = 0 is a local min.

Second derivative test: Checking values of f''(x) on either side of 0,  $f''(-1) = 12(-1)^2 = 12$  and  $f''(1) = 12(1)^2 = 12$ . No change in sign, f'' is positive on either side, so the function is concave up and c = 0 is a local min.

## Example 2: Sketch the function $f'(x) = x^3$

 $f'(x) = 3x^2 = 0$  at x = 0; f''(x) = 6x = 0 at x = 0 also. So, what kind of critical point is c = 0? There are two ways to find out:

First derivative test: Checking values of f'(x) on either side of 0,  $f'(-1) = 3(-1)^2 = 3$ , and  $f'(1) = 3(1)^2 = 3$ . Thus, the function is increasing on either side of c = 0, so c is an inflection point.

Second derivative test: Checking values of f''(x) on either side of 0, f''(-1) = 6(-1) = -6 < 0. f''(1) = 6(1) = 6 > 0. The change in sign indicates the graph is concave down to the left of x = 0 and concave up to the right of it, at x = 0 is an inflection.

## Example 3: Sketch the function $f'(x) = x^{1/3}$

 $f'(x) = \frac{1}{3x^{2/3}}$ . The function is differentiable on its domain, but its derivative cannot equal zero.

However, f'(0) does not exist (division by zero). In the DNE sense, c = 0 is a critical value of the function. (You should see from your graph that the tangent line to the function at x = 0 is a vertical line.)

First derivative test: The function is increasing everywhere, as is easily seen in the graph; algebraically, f'(x) is positive on the domain, observable by the fact that  $x^{2/3}$  is the square of a cube root. Thus, by the first derivative test, the function is everywhere increasing. There is no max or min. Is it an inflection point?

$$f''(x) = \frac{-2}{9x^{5/3}}$$

Second derivative test: f''(x) DNE at x = 0, for the same reason (division by zero). Checking values of f''(x) on either side of 0,  $f''(-1) = (-2/9)(-1)^{5/3} = 2/9 > 0$  (concave up).  $f'(1) = (-2/9)(1)^{5/3} = -2/9 < 0$  (concave down). x = 0 is a point of inflection.