


This unit shows how to use analytical methods of calculus to sketch graphs and interpret them.

Polynomial functions are *differentiable* at all values of their domain (the real numbers). Recall that **critical numbers*** of a function are those $x = c$ where either $f'(c) = 0$ or DNE.



Polynomial functions are ideal for investigating critical numbers of a function, intervals of increase and decrease, extremes, concavity and points of inflection.



Definition: c is a *critical number* of function $f(x)$ if either $f'(c) = 0$ or $f'(c)$ does not exist (DNE).

A function may have a *local extreme* at a critical number c (including a non-differentiable cusp or corner), or it may have an *inflection point*.

Definitions

A function $f(x)$ is (strictly) **increasing** on an interval I if for each a, b in I , when $a < b, f(a) < f(b)$.

A function $f(x)$ is (strictly) **decreasing** on an interval I if for each a, b in I , when $a < b, f(a) > f(b)$.



On an interval where $f(x)$ is increasing, $f'(x) > 0$; on an interval where $f(x)$ is decreasing, $f'(x) < 0$.


$f(x)$ has a **local (relative) maximum** at a if $f(x) \leq f(a)$ for all x in an arbitrarily small interval (called an ϵ -neighborhood) of a .

$f(x)$ has a **local (relative) minimum** at a if $f(x) \geq f(a)$ for all x in an ϵ -neighborhood of a .

[Note: A constant function on I has both a local a max and a local min on that interval.]

Local maximum and minimum values of a function are called **local extremes** of the function.


M is a **global (absolute) maximum** if for every x in I , $f(x) \leq M$; m is a **global (absolute) minimum** if for every x in I , $f(x) \geq m$.




First Derivative Test (FDT) of c for $f(c)$ to be a local max or min

Visually, $f(c)$ is a local max if f is increasing when $x < c$ and decreasing when $x > c$. $f(c)$ is a local min if f is decreasing when $x < c$ and increasing when $x > c$.

Analytically, $f'(x)$ changes sign from positive to negative on either side of c when $f(c)$ is a local max.



And $f'(x)$ changes sign from negative to positive on either side of c when $f(c)$ is a local min.




Thus, the FDT shows whether $f(c)$ is a local max or min by the sign of $f'(x)$ on either side of c .

Second Derivative Test (SDT) of c to for $f(c)$ to be a local max or min

Intervals of the graph where the shape is roughly similar to a cup **concave up**.

Intervals of the graph where the shape is roughly similar to a frown **concave down**.




$f(x)$ is concave up on an interval where $f'(x)$ is increasing because the progress of the slope of the tangent is to increase. Thus, $f''(x) > 0$ on “concave up intervals.”

$f(x)$ is concave down on an interval where $f'(x)$ is decreasing because the progress of the slope of the tangent is to decrease. Thus, $f''(x) < 0$ on “concave down intervals.”

The SDT states that if $f'(c) = 0$ and $f''(c) > 0$, then $f(c)$ is a local min because the graph is concave up there. Likewise, if $f'(c) = 0$ and $f''(c) < 0$, then $f(c)$ is a local max because the graph is concave down there.

An **inflection point** is a point where a graph’s concavity changes.



If $f'(c) = 0$, $f(c)$ could be an inflection point, but not necessarily. Note that $f(x) = x^4$ has both a first and second derivative = 0 at $x = 0$, has an $f(0)$ is a minimum of the function. (See examples below of basic functions and their critical and inflection points.)

To decide what the situation is in the case where $f'(x) = 0$, we do the following:

EITHER

1. Resort to the *first derivative test*, checking values on either side of c to see if $f'(c)$ changes sign.

If it does, we have a local extreme at $x = c$.

If it doesn't, we have an inflection point at $x = c$.

OR

2. Stay with the *second derivative test*, testing values either side of c to see if $f''(x)$ changes sign.

If $f''(x)$ changes sign at c , then c is an *inflection point*, since concavity has changed.

If $f''(x) > 0$ on *both sides* then $x = c$ is a local min.



If $f''(x) < 0$ on *both sides*, then $x = c$ is a local max.

The following examples are illustrative because the functions are so simple to inspect through a sketch.

Example 1

$$f(x) = x^2 \text{ (parabola)}$$

$$f(x) = x^4 \text{ (quartic)}$$



Analyzing the parabola is easy:

FDT: $f'(x) = 2x = 0$ at $x = 0$. $f'(-1) = -2 < 0$; f is decreasing left of zero. $f'(1) = 2 > 0$; f is increasing right of zero. $f(0)$ is a local min.

SDT: $f''(x) = 2 > 0$ for all x , so f is concave up everywhere. Thus, $f(0)$ is a local min, as the graph shows.

Now the quartic:


$f(x) = 4x^3 = 0$ at $x = 0$; $f''(x) = 12x^2 = 0$ at $x = 0$, also. What kind of critical point, then, is $c = 0$?

There are two ways to find out:

FDT: Checking values into $f'(x)$ on either side of 0, $f'(-1) = 4(-1)^3 = -4$ and $f'(1) = 4(1)^3 = 4$. Because f' changes sign, negative to positive, $c = 0$ is a local min.

SDT: Checking values of $f''(x)$ on either side of 0, $f''(-1) = 12(-1)^2 = 12$ and $f''(1) = 12(1)^2 = 12$. No change in sign, f'' is positive on either side, so the function is concave up, and $c = 0$ is a local min, as the graph shows.

Example 2 $f(x) = x^3$ (cubic)




$f(x) = 3x^2 = 0$ at $x = 0$; $f''(x) = 6x = 0$ at $x = 0$ also. So, what kind of critical point is $c = 0$? There are two ways to find out:

FDT: Checking values of $f'(x)$ on either side of 0, $f'(-1) = 3(-1)^2 = 3$, and $f'(1) = 3(1)^2 = 3$. Thus, the function is increasing on either side of $c = 0$, so c is an inflection point, as the graph shows.

SDT: Checking values of $f''(x)$ on either side of 0, $f''(-1) = 6(-1) = -6 < 0$. $f''(1) = 6(1) = 6 > 0$. The change in sign indicates the graph is concave down to the left of $x = 0$ and concave up to the right of it, at $x = 0$ is an inflection.

Example 3

$f(x) = x^{1/3}$ (cube root)



$$f(x) = \frac{1}{3x^{2/3}}$$

We see that $f'(0)$ does not exist (division by zero). In the DNE sense, $c = 0$ is a critical value of the function. (You see from the graph that the tangent line to the function at $x = 0$ is a vertical line.)

FDT: The function is increasing everywhere, as is easily seen in the graph; algebraically, $f(x)$ is *positive* everywhere it is defined, as $x^{2/3}$ is the square of a cube root. Check $f'(-1)$ and $f'(1)$.

Thus, by the first derivative test, the function is everywhere increasing. There is no local max or min.

Is $x = 0$ an inflection point? $f'(x) = -\frac{2}{9x^{5/3}}$

SDT: $f''(x)$ DNE at $x = 0$, for the same reason (division by zero). Checking values of $f''(x)$ on either side of 0:

$$f''(-1) = -\frac{2}{9(-1)^{5/3}} = \frac{2}{9} > 0 \text{ (concave up).}$$

$$f''(1) = -\frac{2}{9} < 0 \text{ (concave down). } x = 0 \text{ is a point of inflection, as the graph shows.}$$

Example 4: (For you to do)

Consider the function $f(x) = |x^2 - 1|$ on \mathbb{R} . Use Desmos to graph it. Look at it for a while. Draw the graph yourself on your paper.

Then write the piecewise function, its first and second derivatives (also piecewise functions) and find critical points. Investigate extremes, intervals of increasing and decreasing, concave up and concave down, any inflection points, and so on.

Interesting conclusion you should be able to draw (and draw with your pencil): A cusp can be a point of inflection.