

## Math 402 - 01 Previous Homework (Spring 2019)

---

```

\newcommand{\aut}{\textrm{Aut}} \newcommand{\inn}{\textrm{Inn}} \newcommand{\sub}{\textrm{Sub}}
\newcommand{\cl}{\textrm{cl}} \newcommand{\join}{\vee} \newcommand{\bigjoin}{\bigvee}
\newcommand{\meet}{\wedge} \newcommand{\bigmeet}{\bigwedge} \newcommand{\normaleq}{\unlhd}
\newcommand{\normal}{\lhd} \newcommand{\union}{\cup} \newcommand{\intersection}{\cap}
\newcommand{\bigunion}{\bigcup} \newcommand{\bigintersection}{\bigcap} \newcommand{\sq}[2][\
]{\sqrt[#1]{#2},} \newcommand{\pbr}[1]{\langle #1\rangle} \newcommand{\ds}{\displaystyle}
\newcommand{\C}{\mathbb{C}} \newcommand{\R}{\mathbb{R}} \newcommand{\Q}{\mathbb{Q}}
\newcommand{\Z}{\mathbb{Z}} \newcommand{\N}{\mathbb{N}} \newcommand{\A}{\mathbb{A}}
\newcommand{\F}{\mathbb{F}} \newcommand{\T}{\mathbb{T}} \newcommand{\ol}[1]{\overline{#1}}
\newcommand{\imp}{\rightarrow} \newcommand{\rimp}{\leftarrow} \newcommand{\pinfty}{1/p^{\infty}}
\newcommand{\power}{\mathcal{P}} \newcommand{\calL}{\mathcal{L}} \newcommand{\calC}{\mathcal{C}}
\newcommand{\calN}{\mathcal{N}} \newcommand{\calB}{\mathcal{B}} \newcommand{\calF}{\mathcal{F}}
\newcommand{\calR}{\mathcal{R}} \newcommand{\calS}{\mathcal{S}} \newcommand{\calU}{\mathcal{U}}
\newcommand{\calT}{\mathcal{T}} \newcommand{\gal}{\textrm{Gal}} \newcommand{\isom}{\approx}
\newcommand{\idl}{\textrm{Idl}} \newcommand{\lub}{\textrm{lub}} \newcommand{\glb}{\textrm{glb}}
\newcommand{\cis}{\textrm{cis}} $

```

### Problem Set 07 (complete) Due: 04/17/2019 Board presentation: 04/26/2019

1. Let  $E$  be a field,  $G$  a finite subgroup of  $\text{Aut}(E)$ ,  $F = E_G$ , and  $L \in \text{Sub}_F(E)$ . Show that  $L^* = \text{Aut}_L(E)$ , and it is a subgroup of  $G$ .
2. Let  $E$  be a field,  $G$  a subgroup of  $\text{Aut}(E)$ , and  $F = E_G$ . Prove that for any  $H, H_1, H_2 \in \text{Sub}(G)$ , and any  $L, L_1, L_2 \in \text{Sub}_F(E)$ 
  - I. If  $H_1 \leq H_2$ , then  $H_2^* \leq H_1^*$ . (i.e.  $\text{Aut}_L^*$  is order reversing)
  - II. If  $L_1 \leq L_2$ , then  $L_2^* \leq L_1^*$ . (i.e.  $\text{Aut}_L^*$  is order reversing)
  - III.  $H \leq H^{**}$  (i.e.  $H \leq \text{Aut}_{H^*}$ )
  - IV.  $L \leq L^{**}$  (i.e.  $L \leq \text{Aut}_{L^*}$ )
3. Let  $E/L/F$  be a field tower.
  - I. Prove that if  $E/F$  is a normal extension then so is  $E/L$ .
  - II. Prove that if  $E/F$  is a Galois extension then so is  $E/L$ .

### Problem Set 06 (complete) Due: 04/12/2019 Board presentation: 04/17/2019

1. Let  $F$  be a field,  $\alpha_1, \dots, \alpha_n$  elements from some extension  $E$  of  $F$ , and  $R$  a commutative ring with unity. If  $\varphi_1, \varphi_2: F(\alpha_1, \dots, \alpha_n) \rightarrow R$  are homomorphisms such that  $\varphi_1(a) = \varphi_2(a)$  for all  $a \in F$  and  $\varphi_1(\alpha_i) = \varphi_2(\alpha_i)$  for  $i = 1, \dots, n$ , then  $\varphi_1 = \varphi_2$ .
2. Let  $f(x) = x^5 - 2 \in \mathbb{Q}[x]$ , and  $E$  the splitting field of  $f(x)$ . Consider the group  $G = \text{Aut}_{\mathbb{Q}}(E)$ .
  - I. What is the order of  $G$ ?
  - II. Is it abelian?
  - III. What are the orders of elements in  $G$ ?
3. Let  $F = \mathbb{F}_p(t)$  be the field of rational functions on  $t$  with coefficients in  $\mathbb{F}_p$ . Consider the polynomial  $f(x) = x^p - t \in F[x]$ .

- I. Show that  $f(x)$  has no root in  $F$ .
- II. Show that the Frobenius endomorphism  $\Phi: F \rightarrow F$  is not surjective.
- III. Show that  $f(x)$  has exactly one root, and that root has multiplicity  $p$ .
- IV. Show that  $f(x)$  is irreducible over  $F$ .

**Problem Set 05** (complete) Due: 03/25/2019 Board presentation: 04/02/2019

1. Let  $F$  be a field and  $f(x), g(x) \in F[x]$ . Prove:
  - I.  $(f(x)+g(x))' = f'(x) + g'(x)$
  - II.  $(f(x)g(x))' = f(x)g'(x) + f'(x)g(x)$
2. Let  $F$  be a field, and  $\varphi: F \rightarrow F$  an endomorphism of  $F$ . Prove that the set  $\{ \{ f \mid \varphi(a) = a \} \}$  is a subfield of  $F$ .
3. How many monic irreducible polynomials of degree 4 are there over  $\mathbb{F}_5$ ?
4. Let  $E$  be a field extension of  $F$ . Prove that  $E$  is an algebraic closure of  $F$  iff  $E$  is minimal with the property that every polynomial  $f(x) \in F[x]$  splits over  $E$ .

**Problem Set 04** (complete) Due: 03/11/2019 Board presentation: 03/25/2019

1. Let  $E/F$  be a field extension. Prove that  $[E:F] = 1$  iff  $E = F$ .
2. Let  $E$  and  $K$  be field extensions of  $F$  and  $\varphi: E \rightarrow K$  an  $F$ -extension homomorphism. Show that  $\varphi$  is a linear transformation of  $F$ -vector spaces.
3. Write  $\sqrt{2}$  as a polynomial expression on  $\alpha = \sqrt{2} + \sqrt{3}$ .
4. Find the minimal polynomial of  $u = (\sqrt{3}^2 + \omega)$  over  $\mathbb{Q}$ .

**Problem Set 03** (complete) Due: 02/18/2019 Board presentation: 02/20/2019

1. Let  $V$  be a vector space and  $B \subseteq V$ . Show that the following are equivalent
  - I.  $B$  is a basis for  $V$ ,
  - II.  $B$  is maximal linearly independent set,
  - III.  $B$  is minimal spanning set.
2. Let  $V$  be a vector space and  $W$  a subspace of  $V$ .
  - I. Prove that  $\dim(W) \leq \dim(V)$ .
  - II. Prove that if  $V$  is finite dimensional and  $\dim(W) = \dim(V)$  then  $W = V$
  - III. Show, with a counterexample, that the finite dimensional hypothesis is necessary in part b.
3. In regards to the *Universal Mapping Property* for vector spaces discussed in class today:
  - I. Complete the proof of it.
  - II. Prove that the set  $\{ \alpha(v) \mid v \in B \}$  is linearly independent in  $W$  iff  $\widehat{\alpha}$  is injective.
  - III. Prove that the set  $\{ \alpha(v) \mid v \in B \}$  is a spanning set for  $W$  iff  $\widehat{\alpha}$  is surjective.
4. Let  $V$  be a vector space over  $F$ , and  $W$  a subspace of  $V$ . Let  $B_1$  be a basis for  $W$  and  $B$  a basis for  $V$  such that  $B_1 \subseteq B$ . Prove that the set  $\{ \{ v+W \mid v \in B - B_1 \} \}$  is a basis for the quotient space  $V/W$ .

**Problem Set 02** (complete) Due: 02/11/2019 Board presentation: 02/18/2019

1. Let  $D$  be a UFD.  $a, b, c \in D$ , and  $f(x) \in D[x]$ .  $a, b$  are said to be "relatively prime" if  $\gcd(a, b)$  is a unit.
  - I. Prove that if  $a, b$  are relatively prime and  $a \mid bc$  then  $a \mid c$ .

- II. Prove that if  $\frac{a}{b}$  is a root of  $f(x)$ , and  $a, b$  are relatively prime, then  $a$  divides the constant term of  $f(x)$  and  $b$  divides the leading term of  $f(x)$ .
2. Let  $D$  be an ED,  $a, b \in D$ , with  $b \neq 0$ . Consider the sequence  $r_0, r_1, r_2, \dots, r_n$  defined recursively as follows:  $r_0 = a, r_1 = b$ , and using the property of an Euclidean Domain, until obtaining a residue  $0$ ,  $\left[ \begin{array}{l} r_0 = a, r_1 = b \\ r_2 = r_1 r_0 + r_2 \text{ and } \delta(r_2) < \delta(r_1) \\ r_3 = r_2 r_1 + r_3 \text{ and } \delta(r_3) < \delta(r_2) \\ \vdots \\ r_{n-3} = r_{n-2} r_{n-1} + r_{n-3} \text{ and } \delta(r_{n-3}) < \delta(r_{n-2}) \\ r_{n-2} = r_{n-1} r_{n-2} + r_n \text{ and } r_n = 0 \end{array} \right]$  Why does the sequence  $r_1, r_2, \dots, r_n$  have to eventually attain the value  $r_n = 0$ ? Prove that the last non-zero entry in the residues list, i.e.  $r_{n-1} \sim \gcd(a, b)$ .
3. Let  $D$  be a PID,  $a, b \in D$ . Let  $d$  be a generator of the ideal  $\langle a \rangle + \langle b \rangle$ . Show that  $d \sim \gcd(a, b)$ .
4. Let  $D$  be an ID,  $a, b \in D$ . Prove that if  $a$  and  $b$  have a least common multiple  $l \in D$ , then  $\frac{ab}{l}$  is a greatest common divisor of  $a$  and  $b$  in  $D$ .
5. (Optional) Let  $\gamma = \mathbb{Z}[\frac{1 + \sqrt{-19}}{2}]$  and consider the subring of  $\mathbb{C}$  given by:  $R = \{a + b\gamma \mid a, b \in \mathbb{Z}\}$  Prove that  $R$  is a PID but not an ED. A detailed proof can be found in Mathematics Magazine, Vol. 46, No. 1 (1973), pp 34-38. If you choose to work on this problem, do not consult this reference, or any other reference. Hand-in only your own work, even if it is only parts of the solution.

**Problem Set 01** (complete) Due: 02/01/2019 Board presentation: 02/08/2019

1. Let  $D$  be an integral domain. Consider the following two properties that  $D$  and a function  $\delta: D \setminus \{0\} \rightarrow \mathbb{N}_0$  may have:
- For any  $a, d \in D$  with  $d \neq 0$ , there are  $q, r \in D$  such that  $a = qd + r$  and ( $r = 0$  or  $\delta(r) < \delta(d)$ )
  - For any  $a, b \in D \setminus \{0\}$ ,  $\delta(a) \leq \delta(ab)$ .
- Prove that if there is a function  $\delta$  satisfying the first condition, then there is a function  $\gamma$  satisfying both of them. Hint: consider  $\gamma$  defined by:  $\gamma(a) = \min_{x \in D \setminus \{0\}} \delta(ax)$
2. Chapter 18, problem 22.
3. Chapter 16, problem 24. Can you weaken the assumption “infinitely many”?
4. Show that an integral domain  $D$  satisfies the ascending chain condition ACC iff every ideal of  $D$  is finitely generated. (Hint: one direction is similar to the proof that every PID satisfies the ACC).

[Home](#)

From:

<http://www2.math.binghamton.edu/> - **Binghamton University Department of Mathematical Sciences**

Permanent link:

[http://www2.math.binghamton.edu/p/people/fer/402ws/spring2019/previous\\_homework](http://www2.math.binghamton.edu/p/people/fer/402ws/spring2019/previous_homework)

Last update: **2019/04/30 19:18**

