2025/09/30 05:42 1/1 Thomas Zaslavsky

## **Thomas Zaslavsky**

## **Biased Graphs and the Associative Law**

## Abstract for the Combinatorics and Number Theory Seminar 2001 April 24

A quasigroup is like a group but without the identity, inverses, or associativity; all that is left is the multiplication table, which is an arbitrary Latin square. This is worth something: one has unique solvability of equations xy=z. Also, an identity can always be found. The crucial missing property is the associative law.

A biased graph is a graph together with a distinguished class of circles (a.k.a. circuits, cycles, polygons) that satisfies a certain combinatorial property. Each quasigroup with m elements gives rise to a kind of biased graph called an m-fold biased expansion of  $K_3$ , the complete graph of order 3. Conversely, every m-fold biased expansion of  $K_3$  is obtained from a quasigroup. Trying to generalize this construction of quasigroups to  $K_n$  fails to be interesting because  $K_4$  implies the associative law, so that a biased expansion of  $K_n$  must come from a group if n > 3.

We explore this fact and possible ways of getting around it. For instance, an m-fold biased expansion of  $C_n$ , the circle of length n, encodes an n-dimensional Latin hypercube, which might be considered the multiplication table of an (n-1)-ary operation. Chords in the circle imply specific associative properties which get stronger and stronger until when all chords have been added (so the graph becomes  $K_n$ ) there is complete associativity, making the operation a group operation. What happens in between is completely unknown.

From:

http://www2.math.binghamton.edu/ - **Department of Mathematics and Statistics, Binghamton University** 

Permanent link:

http://www2.math.binghamton.edu/p/seminars/comb/abstract.200104zas

Last update: 2020/01/29 19:03