

## Tom Head (Binghamton)

---

### Computing Transparently: The Independent Sets in a Graph

---

#### Abstract for the Combinatorics Seminar 2009 October 20

---

I give a procedure for finding the independent sets in an undirected graph by xeroxing onto transparent plastic sheets.

Let an undirected graph having  $n$  vertices and  $m$  edges be given. A list of all the independent subsets of the set of vertices of the graph is constructed by using a xerox machine in a manner that requires the formation of only  $n + m + 1$  successive transparencies. An accompanying list of the counts of the elements in each independent set is then constructed using only  $O(n^2)$  additional transparencies. The list with counts provides a list of all maximum independent sets. This gives an  $O(n^2)$ -step solution for the classical problem of finding the cardinality of a maximal independent set in a graph. The applicability of these procedures is limited, of course, by the increase in the information density on the transparencies when  $n$  is large.

My ultimate purpose here is to give hand tested 'ultra parallel' algorithmic procedures that may prove suitable for realization using future optical technologies.

---

From:

<http://www2.math.binghamton.edu/> - **Binghamton University Department of Mathematical Sciences**

Permanent link:

<http://www2.math.binghamton.edu/p/seminars/comb/abstract.200910hea>

Last update: **2020/01/29 19:03**

