

Steven Simon (Bard)

Regular Polygonal and Prismatic Tverberg-Type Theorems

Abstract for the Combinatorics Seminar 2019 March 26

The celebrated Tverberg Theorem, generalizing Radon's Theorem when $q = 2$, states that any $T(q,d) = (q-1)(d+1)+1$ points in \mathbb{R}^d can be partitioned into q pairwise disjoint sets whose convex hulls have non-empty q -fold intersection. Moreover, this "Tverberg number" $T(q,d)$ is generically tight. We will show that in the absence of a full Tverberg partition for fewer than $T(q,d)$ points, one can nonetheless guarantee a partition of these points into q pairwise disjoint sets so there are q points, one from each of the resulting convex hulls, which form the vertices of a regular q -gon. Analogous results hold for regular prisms when q is composite. As with Tverberg's original theorem, these results can be extended to the continuous setting when q is a prime power, and these likewise admit constrained versions: restrictions on the number of points in each of the disjoint sets (van-Kampen-Flores type results) and the prescription of equal distance from each vertex of the regular polygon to the original set, "balanced" colored versions of a Sobéron variety, and so on.

From:
<http://www2.math.binghamton.edu/> - **Binghamton University Department of Mathematical Sciences**

Permanent link:
<http://www2.math.binghamton.edu/p/seminars/comb/abstract.201903sim>

Last update: **2020/01/29 19:03**

