

Math 447 Spring 2015

Exam 3

April 21, 2015

- Total value 220 points. Each part valued as indicated.
- SHOW YOUR WORK unless otherwise indicated. "NO WORK" may result in "NO POINTS".
- Simplify your answers when possible. Do the arithmetic, remove parentheses, reduce fraction, etc.
- Cross out anything you don't want graded!
- Use the back sides of pages if you need extra space. If you have anything on a back side that you want graded, indicate where it is.

Student: Solution

Section (please circle): 01 (Xu)

Problem #	Possible Points	Points
I	30	
II	14	
III	21	
IV	36	
V	30	
VI	28	
VII	28	
VIII	33	
Total	220	

- I. (30 points.) X has probability mass function (PMF; i.e., $p(x) = P\{X = x\}$) given in the table below

x	-2	0	2	otherwise
$p(x) = P\{X = x\}$	$\frac{1}{6}$	$\frac{2}{3}$	$\frac{1}{6}$	0

- (1) (8 points) Give the distribution function (CDF) of X .
I.e., $F(x) = P\{X \leq x\} = ?$

$$F(x) = \begin{cases} 0 & x < -2 \\ \frac{1}{6} & -2 \leq x < 0 \\ \frac{5}{6} & 0 \leq x < 2 \\ 1 & x \geq 2 \end{cases}$$

- (2) (6 points) $EX^4 = ?$

$$= (-2)^4 \times \frac{1}{6} + 0^4 \times \frac{2}{3} + 2^4 \times \frac{1}{6} = \frac{16}{6} + \frac{16}{6} = \frac{32}{6} = \frac{16}{3}$$

- (3) (6 points) $\text{Var } X = ?$

$$EX = (-2) \times \frac{1}{6} + 0 \times \frac{2}{3} + 2 \times \frac{1}{6} = 0$$

$$EX^2 = (-2)^2 \times \frac{1}{6} + 0^2 \times \frac{2}{3} + 2^2 \times \frac{1}{6} = \frac{4}{3} \Rightarrow \text{Var}(X) = EX^2 - (EX)^2 = \frac{4}{3} - 0^2 = \frac{4}{3}$$

- (4) (6 points) Define $Y = \frac{1}{X+1}$. Give the moment generating function (MGF) of Y , $m_Y(t) = ?$.

$$M_Y(t) = Ee^{tY} = Ee^{\frac{t}{X+1}} = e^{\frac{t}{-2+1}} p(X=-2) + e^{\frac{t}{0+1}} p(X=0) + e^{\frac{t}{2+1}} p(X=2)$$

$$= \frac{1}{6} e^{-t} + \frac{2}{3} e^t + \frac{1}{6} e^{\frac{t}{3}}$$

- (5) (4 points) Find the q th quantile of X with $q = \frac{1}{4}$. (No work need be shown.)

2

0

II. (14 points.) $\{X_1, \dots, X_n\}$ is an independent collection of random variables having finite mean μ and finite variance σ^2 . Let $S = \sum_{k=1}^3 kX_k$. Your answers to the questions below will involve μ and σ .

(1) (4 points.) $E(S) = ?$

$$E(S) = E\left(\sum_{k=1}^3 kX_k\right) = \sum_{k=1}^3 kE(X_k) = \sum_{k=1}^3 k\mu = \mu(1+2+3) = 6\mu$$

(independent)

(2) (4 points) $Var(S) = ?$

$$Var(S) = Var\left(\sum_{k=1}^3 kX_k\right) = \sum_{k=1}^3 Var(kX_k) - 2 \sum_{k=1}^3 \sum_{j=2}^3 \underbrace{Cov(kX_k, jX_j)}_{\rho}$$

$$= \sum_{k=1}^3 k^2 Var(X_k) - 0 = \sum_{k=1}^3 k^2 \cdot \sigma^2 = (1+2^2+3^2)\sigma^2 = 14\sigma^2$$

(3) (6 points) If $Y = X_1 + 2X_2 - 3$ and $Z = X_1 - 2X_2 + X_3 + 4$, then $Cov(Y, Z) = ?$

$$Cov(Y, Z) = Cov(X_1 + 2X_2 - 3, X_1 - 2X_2 + X_3 + 4)$$

$$= Cov(X_1 + 2X_2, X_1 - 2X_2 + X_3)$$

$$= Cov(X_1, X_1) - 2Cov(X_1, X_2) + Cov(X_1, X_3)$$

$$+ (2)Cov(X_2, X_1) + (2)(-2)Cov(X_2, X_2)$$

$$+ 2Cov(X_2, X_3)$$

$$= Var(X_1) - 4Var(X_2)$$

$$= \sigma^2 - 4\sigma^2$$

$$= -3\sigma^2$$

III. (21 points.) X has probability density function (PDF)

$$f_X(x) = \begin{cases} \frac{3x^2}{8} & 0 < x < 2 \\ 0 & \text{otherwise.} \end{cases}$$

(1) (6 points) Give the distribution function (CDF) of X .

I.e., $F(x) = P\{X \leq x\} = ?$ (Be sure to give all pieces of this function. Otherwise you will lose points.)

$$F(x) = \int_{-\infty}^x f_X(t) dt = \begin{cases} 0, & x \leq 0 \\ \int_0^x \frac{3}{8} t^2 dt, & 0 < x < 2 \\ 1, & x \geq 2 \end{cases} = \begin{cases} 0, & x \leq 0 \\ \frac{1}{8} x^3, & 0 < x < 2 \\ 1, & x \geq 2 \end{cases}$$

(2) (6 points) $EX = ?$

$$\int_{-\infty}^{\infty} x f_X(x) dx = \int_0^2 \frac{3}{8} x^3 dx = \frac{3}{32} x^4 \Big|_0^2$$
$$= \frac{3}{32} \times 16 = \frac{3}{2}$$

(3) (3 points) $P\{X = 1/3\} = ?$ \emptyset since X is continuous.

(4) (6 points) $P\{X^2 < \frac{1}{4}\} = ?$

$$P\left(-\frac{1}{2} < X < \frac{1}{2}\right) = P\left(0 < X < \frac{1}{2}\right)$$
$$= F\left(\frac{1}{2}\right) - F(0) = \frac{1}{8} \times \left(\frac{1}{2}\right)^3 = \frac{1}{64}$$

IV. (36 points.) X has distribution function (CDF)

$$F_X(x) = \begin{cases} 0 & x < 0 \\ \frac{1+x^2}{10} & 0 \leq x < 2 \\ 1 & 2 \leq x \end{cases}$$

- (1) (5 points) Does this distribution function (CDF) have any jumps? If yes, identify all the jump points and the associated probabilities (i.e. $P(X = x) = ?$).

$$P(X=0) = F(0) - F(0^-) = \frac{1+0^2}{10} - 0 = \frac{1}{10}$$

$$P(X=2) = F(2) - F(2^-) = 1 - \frac{1+2^2}{10} = \frac{1}{2}$$

- (2) (5 points) $P\{-\pi < X \leq \frac{1}{2}\} = ?$ $F(\frac{1}{2}) - F(-\pi) = \frac{1+(\frac{1}{2})^2}{10} - 0 = \frac{1}{8}$

- (3) (5 points) $P\{0 \leq x \leq 2\} = ?$ $F(2) - F(0^-) = 1 - 0 = 1$

- (4) (3 points) A is the set $\{-1, 0, \frac{1}{\sqrt{2}}, 2\}$. $P\{X \notin A\} = ?$

$$= 1 - P(X \in A) = 1 - (P(X=0) + P(X=2) + 0 + 0)$$

$$= 1 - \left(\frac{1}{10} + \frac{1}{2}\right) = \frac{2}{5}$$

(5) (2 points) Find $F'(x) = ?$

$$\begin{cases} \frac{x}{5}, & 0 < x < 2 \\ 0, & \text{elsewhere.} \end{cases}$$

(6) (8 points) $EX = ?$

$$\begin{aligned} & \int_{-\infty}^{+\infty} x F'(x) dx + 0 \times p(x=0) + 2 p(x=2) \\ &= \int_0^2 \frac{x^2}{5} dx + 2 \times \frac{1}{2} \\ &= \frac{x^3}{15} \Big|_0^2 + 1 = \frac{23}{15} \end{aligned}$$

(7) (8 points) $Var(X) = ?$ (give a summation of reduced fractions as your final answer, no need to simplify.)

$$\begin{aligned} EX^2 &= \int_{-\infty}^{+\infty} x^2 F'(x) dx + 0^2 p(x=0) + 2^2 p(x=2) \\ &= \int_0^2 \frac{x^3}{5} dx + 4 \times \frac{1}{2} = \frac{1}{20} x^4 \Big|_0^2 + 2 = \frac{14}{5} \end{aligned}$$

$$Var(X) = EX^2 - (EX)^2 = \frac{14}{5} - \left(\frac{23}{15}\right)^2 = \frac{101}{225}$$

V. (30 points.) X is normally distributed with mean $\mu = 3$ and variance $\sigma^2 = 36$.

(1) (5 points) $P\{3 \leq X \leq 15\} = ?$ $P\left(\frac{3-3}{6} \leq Z \leq \frac{15-3}{6}\right) = P(0 \leq Z \leq 2)$

$$= \Phi(2) - \Phi(0) = 0.9772 - 0.5 = 0.4772$$

(2) (5 points) $P\{|X - 6| \leq 9\} = ?$ $P(-3 \leq X \leq 15) = P\left(\frac{-3-3}{6} \leq Z \leq \frac{15-3}{6}\right)$

$$= P(-1 \leq Z \leq 2) = \Phi(2) - \Phi(-1) = \Phi(2) - (1 - \Phi(1))$$

$$= \Phi(2) + \Phi(1) - 1 = 0.9772 + 0.8413 - 1 = 0.8185$$

(3) (5 points) Find out a value for x_0 so that $P\{X \geq x_0\} = 0.99$

$$P(X \geq x_0) = 0.99 \Rightarrow P\left(Z \geq \frac{x_0 - 3}{6}\right) = 0.99$$

$$\Rightarrow P\left(Z \leq -\frac{x_0 - 3}{6}\right) = 0.99 \Rightarrow -\frac{x_0 - 3}{6} = Z_{0.99} = 2.33$$

$$\Rightarrow x_0 = 3 - 2.33 \times 6 = \boxed{-10.98}$$

(4) (5 points) $P\{X^3 + 27 \geq 0\} = ?$

$$= P(X^3 \geq -27) = P(X \geq -3) = P\left(Z \geq \frac{-3-3}{6}\right) = P(Z \geq -1)$$

$$= P(Z \leq 1) = \Phi(1) = 0.8413$$

(5) (5 points) Give the moment generating function (MGF) for X , $m_X(t) = ?$

$$M_X = e^{3t + \frac{36}{2}t^2} = e^{3t + 18t^2}$$

(6) (5 points) Let $Y = 5X + 10$. Give the probability density function (PDF) for Y .

I.e., $f_Y(x) = ?$

$$EY = 5EX + 10 = 5 \times 3 + 10 = 25$$

$$\text{Var}(Y) = \text{Var}(5X + 10) = 5^2 \text{Var}(X) = 5^2 \cdot 6^2 = 30^2$$

$$Y \sim N(25, 30^2) \quad f_Y(y) = \frac{1}{\sqrt{2\pi} \cdot 30} e^{-\frac{(y-25)^2}{2 \times 30^2}} \quad -\infty < y < +\infty$$

VI. (28 points.) X has a cumulative distribution function (CDF). (If you don't pay attention to the difference between CDF and PDF, you can say goodbye to the following 33 pts)

$$F_X(x) = \begin{cases} 1 - e^{-5x} & x > 0 \\ 0 & x \leq 0 \end{cases}$$

$$X \sim \exp\left(\frac{1}{5}\right)$$

$$\Rightarrow EX = \frac{1}{5}, \text{Var}(X) = \frac{1}{5^2}$$

(1) (4 points) $E(5X + 1) = ?$

$$= 5EX + 1 = 5 \times \frac{1}{5} + 1 = 2$$

(2) (4 points) $\text{Var}(5X + 1) = ?$

$$\text{Var}(5X + 1) = 5^2 \text{Var}(X) = 5^2 \times \frac{1}{5^2} = 1$$

(3) (5 points) Give the moment generating function (MGF) of X , $m_X(t) = ?$

$$M_X(t) = \frac{1}{1 - \frac{1}{5}t} \quad \text{for } (t < 5)$$

(4) (5 points) $E[(5X)^{20}] = ?$

$$E[(5X)^{20}] = 5^{20} EX^{20} = 5^{20} \cdot 20! \left(\frac{1}{5}\right)^{20} = 20!$$

(5) (5 points) $P\left\{X \leq \frac{1}{10} \mid X \geq \frac{1}{20}\right\} = ?$

$$\begin{aligned} &= \frac{P(X \leq \frac{1}{10}, X \geq \frac{1}{20})}{P(X \geq \frac{1}{20})} = \frac{P(\frac{1}{20} \leq X \leq \frac{1}{10})}{P(X \geq \frac{1}{20})} = \frac{F_X(\frac{1}{10}) - F_X(\frac{1}{20})}{1 - F_X(\frac{1}{20})} \\ &= \frac{(1 - e^{-5 \times \frac{1}{10}}) - (1 - e^{-5 \times \frac{1}{20}})}{1 - (1 - e^{-5 \times \frac{1}{20}})} = \frac{e^{-\frac{1}{2}} - e^{-\frac{1}{4}}}{e^{-\frac{1}{4}}} = \boxed{1 - e^{-\frac{1}{4}}} \end{aligned}$$

(6) (5 points) Let $Y = 10X$. Give the distribution function (CDF) of Y . I.e., $F_Y(y) = ?$

Moreover, based on $F_Y(y)$, tell what distribution Y has (and what is/are the parameter(s)).

$$\begin{aligned} F_Y(y) &= P(Y \leq y) = P(10X \leq y) = P\left(X \leq \frac{y}{10}\right) = F_X\left(\frac{y}{10}\right) \\ &= \begin{cases} 1 - e^{-5 \times \frac{y}{10}}, & \frac{y}{10} > 0 \\ 0, & \frac{y}{10} \leq 0 \end{cases} = \begin{cases} 1 - e^{-\frac{y}{2}}, & y > 0 \\ 0, & y \leq 0 \end{cases} \end{aligned}$$

Hence, $Y \sim \text{exponential}(\beta)$ with $\beta = 2$.

VII. (28 points.) X and Y have joint probability mass function given in the table below.

$y \backslash x$	1	2	otherwise
0	$\frac{1}{10}$	$\frac{2}{10}$	0
1	$\frac{3}{10}$	$\frac{4}{10}$	0
otherwise	0	0	0

(1) (5 points) $E(XY) = ?$

$$E(XY) = (1 \times 0) p(1,0) + (1 \times 1) p(1,1) + (2 \times 0) p(2,0) + (2 \times 1) p(2,1)$$

$$= 1 \times \frac{3}{10} + 2 \times \frac{4}{10} = \frac{11}{10}$$

(2) (5 points) $Cov(X, Y) = ?$

$$E(XY) - (E(X)E(Y)) = \frac{11}{10} - \frac{8}{5} \times \frac{7}{10} = \boxed{-\frac{1}{50}}$$

$$E(X) = 1 \times (\frac{1}{10} + \frac{3}{10}) + 2 \times (\frac{2}{10} + \frac{4}{10}) = \frac{8}{5}$$

$$E(Y) = 0 \times (\frac{1}{10} + \frac{2}{10}) + 1 \times (\frac{3}{10} + \frac{4}{10}) = \frac{7}{10}$$

(3) (5 points) Are X and Y independent of each other? (give your arguments. no argument, no point.)

Solution I: $P(X=1, Y=0) = \frac{1}{10} \neq P(X=1)P(Y=0) = \frac{4}{10} \times \frac{3}{10} = \frac{3}{25}$

Solution II: $Cov(X, Y) = -\frac{1}{50} \neq 0$

$\Rightarrow X$ and Y are NOT independent

(4) (8 points) Give $P\{Y = y \mid X = x\}$ for all values of x and y . (You need to organize them as a "table".)

$Y \backslash X$	1	2
0	$\frac{1}{4}$	$\frac{1}{3}$
1	$\frac{3}{4}$	$\frac{2}{3}$

(5) (5 points) $E\{Y \mid X = 1\} = ?$

$$E(Y \mid X=1) = 0 \times p(Y=0 \mid X=1) + 1 \times p(Y=1 \mid X=1)$$

$$= 0 + 1 \times \frac{3}{4} = \frac{3}{4}$$

VIII. (33 points.) X and Y have joint probability density function (PDF)

$$f(x, y) = \begin{cases} \frac{3}{4}y & 0 < x < 2, 0 < y < 2, x + y < 2 \\ 0 & \text{otherwise.} \end{cases}$$

- (1) (5 points) Give the (marginal) probability density (marginal PDF) $f_X(x)$ of X . (Be sure to specify the regions of x .)

$$f_X(x) = \int_{-\infty}^{+\infty} f(x, y) dy = \begin{cases} \int_0^{2-x} \frac{3}{4}y dy = \frac{3}{8}(2-x)^2, & x \in (0, 2) \\ 0, & x \notin (0, 2) \end{cases}$$

- (2) (5 points) Give the (marginal) probability density (marginal PDF) $f_Y(y)$ of Y . (Be sure to specify the regions of y .)

$$f_Y(y) = \int_{-\infty}^{+\infty} f(x, y) dx = \begin{cases} \int_0^{2-y} \frac{3}{4}y dx = \frac{3}{4}y \cdot x \Big|_0^{2-y} = \frac{3}{4}y(2-y), & 0 < y < 2 \\ 0, & y \notin (0, 2) \end{cases}$$

- (3) (4 points) Are X and Y independent of each other? (give your arguments. no argument, no points!)

No, X and Y are NOT independent


because $f(x, y) \neq f_X(x) f_Y(y)$.

- (4) (5 points) $P(X > Y) = ?$ (Leave your answer as an integral. Make sure your limits of integration are correct.)

Solution ①

$$P(X > Y) = \int_0^1 \left(\int_y^{2-y} \frac{3}{4}y dx \right) dy = \int_0^1 \frac{3}{4}y(2-2y) dy$$

$$= \int_0^1 \frac{3}{2}(y-y^2) dy = \left(\frac{3}{4}y^2 - \frac{1}{2}y^3 \right) \Big|_0^1 = \frac{3}{4} - \frac{1}{2} = \boxed{\frac{1}{4}}$$



Solution ② ← (dy) first and then (dx)

$$P(X > Y) = \int_0^1 \left(\int_0^x \frac{3}{4}y dy \right) dx + \int_1^2 \left(\int_0^{2-x} \frac{3}{4}y dy \right) dx$$

$$= \int_0^1 \frac{3}{8}x^2 dx + \int_1^2 \frac{3}{8}(2-x)^2 dx$$

$$= \frac{1}{8}x^3 \Big|_0^1 + \frac{1}{8}(x-2)^3 \Big|_1^2 = \boxed{\frac{1}{4}}$$

- (5) (6 points) Give the conditional probability density function (conditional PDF) of Y given that $X = x$. I.e., $f_{Y|X}(y|x) = ?$ (Be sure to specify the region of (x, y) in which your result holds.)

$$f_{Y|X}(y|x) = \frac{f(x, y)}{f_X(x)} = \begin{cases} \frac{\frac{3}{4}y}{\frac{3}{8}(2-x)^2} = \frac{2y}{(2-x)^2}, & 0 < y < 2-x \\ 0, & y \notin (0, 2-x) \end{cases} \quad x \in (0, 2)$$

undefined, $x \notin (0, 2)$.

- (6) (4 points) $E\{Y | X = \frac{1}{2}\} = ?$

$$\begin{aligned} E(Y|X = \frac{1}{2}) &= \int_{-\infty}^{+\infty} y f_{Y|X=\frac{1}{2}}(y) dy = \int_0^{2-\frac{1}{2}} \frac{2y^2}{(2-\frac{1}{2})^2} dy \\ &= \int_0^{\frac{3}{2}} \frac{8}{9} y^2 dy = \frac{8}{27} y^3 \Big|_0^{\frac{3}{2}} = \frac{8}{27} \times \frac{27}{8} = 1 \end{aligned}$$

- (7) (4 points) $E\{(Y+1)X | X = \frac{1}{2}\} = ?$

$$\begin{aligned} E((Y+1)X | X = \frac{1}{2}) &= E\left(\frac{Y+1}{2} | X = \frac{1}{2}\right) \\ &= \frac{1}{2} E(Y | X = \frac{1}{2}) + \frac{1}{2} \\ &= \frac{1}{2} \times 1 + \frac{1}{2} \\ &= 1 \end{aligned}$$

