


[illegible]



**Problem 1.** [15 pts] Compute the values of the following trigonometric functions at the given points:

$$(a) \cos\left(\frac{7\pi}{6}\right) = -\frac{\sqrt{3}}{2}$$



$$(b) \tan\left(\frac{9\pi}{4}\right) = 1$$



$$\begin{aligned} \sin\left(\frac{9\pi}{4}\right) &= \sin\left(\frac{\pi}{4}\right) = \frac{1}{\sqrt{2}} \\ \cos\left(\frac{9\pi}{4}\right) &= \cos\left(\frac{\pi}{4}\right) = \frac{1}{\sqrt{2}} \end{aligned}$$



$$(c) \sec\left(-\frac{2\pi}{3}\right) =$$

$$= \frac{1}{\cos\left(\frac{4\pi}{3}\right)} = -2$$

$$\begin{aligned} \cos\left(-\frac{2\pi}{3}\right) &= \cos\left(\frac{4\pi}{3}\right) \\ &= -\frac{1}{2} \end{aligned}$$



**Problem 2.** [10 pts] Expand as a sum or difference of logarithms and simplify as much as possible:

$$\log_2 \left( \frac{x^2 + 2x + 8}{16(x^2 + 4)(x + 2)} \right) = \dots$$

$$\dots = \log_2(x^2 + 2x + 8) - [\log_2(16) + \log_2(x^2 + 4) + \log_2(x + 2)]$$

$$= \log_2(x^2 + 2x + 8) - \log_2(16) - \log_2(x^2 + 4) - \log_2(x + 2)$$

$$= \log_2(x^2 + 2x + 8) - 4 - \log_2(x^2 + 4) - \log_2(x + 2)$$

**Problem 3.** [15 pts] Find the exact value of each expression.


$$(a) \arctan(-1) = -\frac{\pi}{4}$$

$$(b) \arcsin\left(\frac{1}{2}\right) = \frac{\pi}{6}$$

$$(c) e^{\ln(27)} = 27$$

$$(d) \arctan(\tan(\pi)) = \arctan(0) = 0$$

$$(e) \log_{25}\left(\frac{1}{5}\right) = -\frac{1}{2} \quad \text{because } 25^{-\frac{1}{2}} = \frac{1}{\sqrt{25}} = \frac{1}{5}$$



**Problem 4.** [5 pts] Rewrite the expression as an algebraic expression in  $x$ :

$$\tan(\arccos(x)) = \frac{\sqrt{1-x^2}}{x}$$


Method 1  $\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)} = \frac{\sqrt{1-\cos^2(\theta)}}{\cos(\theta)} \quad \text{for } \theta \in (0, \pi)$

$$\tan(\arccos(x)) = \frac{\sqrt{1-\cos^2(\arccos(x))}}{\cos(\arccos(x))} = \frac{\sqrt{1-x^2}}{x}$$

Method 2

$$\theta = \arccos(x)$$

$$\tan(\arccos(x)) =$$



$$\tan(\theta) = \frac{\sqrt{1-x^2}}{x}$$

**Problem 5.** [15 pts] Solve each equation below:

(a)  $\ln(3x + 4) = 42$

$$\begin{aligned} 3x + 4 &= e^{42} \\ 3x &= e^{42} - 4 \\ x &= \frac{e^{42} - 4}{3} \end{aligned}$$

(b)  $e^{x^2 - 1} = 8$

$$x^2 - 1 = \ln(e^{x^2 - 1}) = \ln(8)$$

$$x^2 = \ln(8) + 1$$

$$x = \sqrt{\ln(8) + 1} \quad \text{or} \quad -\sqrt{\ln(8) + 1}$$

(Note:  $\ln(8) + 1 > 0$  because  $8 > 1$ .)


(c)  $6 \tan^2(x) - 2 = 0$  in the interval  $[0, 2\pi]$ .

$$2(\tan^2(x) - 1) = 0$$

$$2(\tan(x) - 1)(\tan(x) + 1) = 0$$

$$\tan(x) = 1 \quad \text{or} \quad \tan(x) = -1$$

$$x = \frac{\pi}{4} \quad \text{or} \quad \frac{5\pi}{4} \quad x = \frac{3\pi}{4} \quad \text{or} \quad \frac{7\pi}{4}$$



**Problem 6.** [10 pts] A sum of \$8,000 is invested in a bank that pays interest *continuously* at an annual rate of 5%. If the sum is left in the bank to accumulate interest, how many years will it take for the account to reach \$32,000? [Your answer should be written in a form that could be entered into a calculator for an approximation. Do not attempt to simplify.]

$$P(t) = 8000 e^{.05t}$$

(t measured in years).

$$32000 = 8000 e^{.05t}$$


$$e^{.05t} = 4$$

$$.05t = \ln(4) \quad (\text{or } \log(4))$$


$$t = \frac{\ln(4)}{.05} = 20 \ln(4) \text{ years.}$$

**Problem 7.** [10 pts] Sketch graphs of the following functions. Be sure to label the  $x$ - and  $y$ -intercepts and all asymptotes and to put a *scale* on your axes!

(a)  $-\ln(1-x)$




(b)  $1 + \sin(x - \frac{\pi}{2})$  over the interval  $[-\pi, \pi]$



**Problem 8.** [10 pts]


- (a) On the axes below, sketch a function  $f(x)$  with the following properties. (If any are *impossible*, say so and explain briefly why.)

$$\begin{array}{lll}
 f(-1) = 3 & \lim_{x \rightarrow -1} f(x) = 2 & \\
 f(1) = -1 & \lim_{x \rightarrow 1^-} f(x) = -1 & \lim_{x \rightarrow 1^+} f(x) = 2 \\
 f(2) = 1 & f(x) \text{ is continuous at } 2 & 
 \end{array}$$



- (b) On the axes below, sketch a function  $g(x)$  with the following properties. (If any are *impossible*, say so and explain briefly why.)

$$\begin{array}{lll}
 g(1) = 2 & \lim_{x \rightarrow 1^-} g(x) = \infty & \lim_{x \rightarrow 1^+} g(x) = 2 \\
 g(2) \text{ is undefined} & \lim_{x \rightarrow 2^-} g(x) = 0 & \lim_{x \rightarrow 2^+} g(x) = 0
 \end{array}$$





**Problem 9.** [15 pts] Compute the following limits.

$$(a) \lim_{x \rightarrow 6} \frac{\sqrt{x+3} - 3}{x-6} =$$

$$= \lim_{x \rightarrow 6} \left( \frac{\sqrt{x+3} - 3}{x-6} \right) \left( \frac{\sqrt{x+3} + 3}{\sqrt{x+3} + 3} \right) \quad \text{because } \frac{\sqrt{x+3} + 3}{\sqrt{x+3} + 3} = 1$$

$$= \lim_{x \rightarrow 6} \frac{(x+3) - 9}{(x-6)(\sqrt{x+3} + 3)}$$

$$= \lim_{x \rightarrow 6} \frac{x-6}{(x-6)(\sqrt{x+3} + 3)}$$

$$= \lim_{x \rightarrow 6} \frac{1}{\sqrt{x+3} + 3} \quad \left( \text{because } \frac{x-6}{x-6} = 1 \text{ near } x=6 \right)$$

(limits are local)

$$= \frac{\lim_{x \rightarrow 6} 1}{\lim_{x \rightarrow 6} \sqrt{x+3} + \lim_{x \rightarrow 6} 3} \quad \text{(quotient theorem)}$$

$$= \frac{1}{\sqrt{9} + 3} = \frac{1}{6}$$

$$(b) \lim_{x \rightarrow 5} \frac{x^2 - 3x - 10}{x-5} =$$

$$= \lim_{x \rightarrow 5} \frac{(x-5)(x+2)}{x-5}$$

$$= \lim_{x \rightarrow 5} x+2 \quad \left( \text{because } \frac{x-5}{x-5} = 1 \text{ near } 5 \right)$$

(limits are local)

$$= \lim_{x \rightarrow 5} x + \lim_{x \rightarrow 5} 2 \quad \text{(sum theorem)}$$

$$= 5 + 2 = 7$$

$$(c) \lim_{x \rightarrow 5^-} \frac{25 - x^2}{|x-5|} =$$

$$= \lim_{x \rightarrow 5} \frac{25 - x^2}{-(x-5)} \quad \left( \text{because } |x-5| = -(x-5) \text{ when } x < 5 \right)$$

$$= \lim_{x \rightarrow 5} \frac{(5-x)(5+x)}{5-x}$$

$$= \lim_{x \rightarrow 5} 5+x \quad \left( \text{because } \frac{5-x}{5-x} = 1 \text{ when } x \text{ is near } 5 \right)$$

(limits are local)

$$= \lim_{x \rightarrow 5} 5 + \lim_{x \rightarrow 5} x$$

$$= 5 + 5 = 10$$