

TeX code compiled with `\documentclass{beamer}` using the Amsterdam theme.

There are four png images needed to compile slides:

contour1.png

contour2.png

astroid.png

lemniscate.png

```
\begin{document} \begin{frame} \large Draw a graph of  $x = \sin y$  and find the slope of the line tangent to the graph at the point  $(0, \pi)$ . \vskip 45pt Find  $dx/dy$  and  $dy/dx$  if  $y/\sec(x) = 6x\tan(y)$ . \end{frame}
\begin{frame} \large Find  $dy/dx$  by implicit differentiation. \vskip 15pt \begin{columns} \begin{column}{0.5\textwidth} \begin{enumerate} \item[\bf a)]  $x^4 + y^3 = 1$  \vskip 30pt \item[\bf b)]  $7x^2 + 5xy - y^2 = 6$  \vskip 30pt \item[\bf c)]  $x^7(x + y) = y^2(4x + y)$  \end{enumerate} \end{column} \begin{column}{0.5\textwidth} \begin{enumerate} \item[\bf d)]  $\cos(x)\sin(y) = 2$  \vskip 30pt \item[\bf e)]  $5y\sin(x^2) = 9x\sin(y^2)$  \vskip 30pt \item[\bf f)]  $\sqrt{7x+y} = 6 + x^2y^2$  \end{enumerate} \end{column} \end{columns} \end{frame}
\begin{frame} \large Explain (without calculating) why the two following equations will yield the same formula for  $dy/dx$ . Does this mean that the two graphs will have exactly the same tangent lines?  $x^3y + y^2 + y = 1$   $x^3y + y^2 + y = -1$  \begin{columns} \begin{column}{0.5\textwidth} \includegraphics[height=5cm]{contour1.png} \end{column} \begin{column}{0.5\textwidth} \includegraphics[height=5cm]{contour2.png} \end{column} \end{columns} \end{frame}
\begin{frame} \large Find an equation of the tangent line to the ellipse  $9x^2 + xy + 9y^2 = 19$  at the point  $(1, 1)$ . \vskip 20pt Find an equation of the tangent line to the astroid  $x^{2/3} + y^{2/3} = 4$  at  $(-\sqrt[3]{3}, 1)$ . \begin{center} \includegraphics[height=4cm]{astroid.png} \end{center} \end{frame}
\begin{frame} \large Find the points on the lemniscate  $8(x^2 + y^2)^2 = 25(x^2 - y^2)$  where the tangent is horizontal. \begin{figure}[htp] \centering \includegraphics[height=4cm]{lemniscate.png} \end{figure} \end{frame}
\begin{frame} \large If  $f(x) + x^2[f(x)]^3 = 10$  and  $f(1) = 2$ , find  $f'(1)$ . \vskip 50pt Find  $dx/dy$  and  $dy/dx$  and  $dz/dx$  if  $y/\sec(z) = 6x\tan(y)$ . \end{frame}
\begin{frame} \large Find  $y''$  by implicit differentiation.  $4x^2 + y^2 = 9$  \end{frame}
\begin{frame} \large When we introduced the Power Rule, we explained it for  $y = x^n$  when  $n$  is a nonnegative integer, and we promised that later we'd explain it when  $n$  is a rational and/or negative number. The moment has come. In the following, you should use the Power Rule only for  $n$  a nonnegative integer to prove it the Power Rule for all rational numbers. \begin{enumerate} \item Warm-up: write  $y = x^{\frac{2}{3}}$  as  $y^3 = x^2$ . Then use Implicit Differentiation to show  $y' = \frac{2}{3}x^{-\frac{1}{3}}$ . \item Let  $y = x^{\frac{p}{q}}$ , where  $p$  and  $q$  are positive integers. Use the same method as the previous problem to show  $y' = \frac{p}{q}x^{\frac{p-1}{q}}$ . \item Warm-up: write  $y = x^{-1}$  as  $xy = 1$ . Then use Implicit Differentiation to show  $y' = -x^{-2}$ . \item Let  $y = x^{-a}$ , where  $a$  is a positive rational number. Use the same method as the previous problem to show  $y' = -ax^{-a-1}$ . \end{enumerate} \end{frame}
\begin{frame} \large When you solve for  $y''$  in an implicit differentiation problem, you have to solve a quadratic equation \begin{enumerate} \item always \item sometimes \item never \end{enumerate} \end{frame}
\begin{frame} \large Find equations of both the tangent lines to the ellipse  $x^2 + 9y^2 = 81$  that pass through the point  $(27, 3)$ . \end{frame}
\begin{frame} \large The Thin Lens Equation in optics relates the focal length  $f$  of a lens, the distance  $a$  from an object to the lens, and the distance  $b$  from the object's image to the lens. The equation is  $\frac{1}{a} + \frac{1}{b} = \frac{1}{f}$ . Let's say you have a lens with focal length 10 cm. \begin{itemize} \item Which of the following derivatives describes the rate at which the position of the image changes as you move the object? \centerline{  $\frac{db}{da}$  } \item If the object is 20 centimeters 

```

from the lens and moving away from the lens, where is the object's image and in what direction is it moving?
\end{itemize} \end{frame} \end{document}

From:
<http://www2.math.binghamton.edu/> - **Department of Mathematics and Statistics, Binghamton University**

Permanent link:
http://www2.math.binghamton.edu/p/calculus/resources/calculus_flipped_resources/derivatives/2.6_implicit_differentiation_tex

Last update: **2015/08/29 03:34**