2025/09/16 14:00 1/1 Matthias Beck

## **Matthias Beck**

## On Sperner's Inequality and Its Generalizations

Abstract for the Combinatorics and Number Theory Seminar 2001 December 6

Sperner's inequality concerns the set P(S) consisting of all subsets of a finite set S, partially ordered by subset inclusion. A *chain* is a linearly ordered subset of P(S). In 1928, Sperner found a bound (in terms of the size of S) for the size of an antichain, that is, a subset of P(S) in which there are no nontrivial chains. His result was generalized in two different directions: About 25 years later, Erdös extended Sperner's inequality to r-systems, that is, subsets of P(S) in which chains contain at most r elements. Roughly another 25 years later, Meshalkin proved a Sperner-like inequality for a certain family of compositions (ordered partitions) of S. It's been about 25 years since Meshalkin's theorem, and hence about time for another extension. Xueqin Wang and I unify Erdös's and Meshalkin's inequalities and generalize them to a less restrictive family of compositions of S. As with the known theorems, our result has analogues in the poset of subspaces of a vector space.

From:

http://www2.math.binghamton.edu/ - **Department of Mathematics and Statistics, Binghamton University** 

Permanent link:

http://www2.math.binghamton.edu/p/seminars/comb/abstract.200012beck

Last update: 2020/01/29 19:03