

Béla Bajnok

Spherical Designs and the Independence Number of a Subset of an Abelian Group

Abstract for the Combinatorics and Number Theory Seminar 2002 March 13

A finite subset X on the unit sphere S^d in \mathbb{R}^{d+1} is a *spherical t -design* if for every polynomial $f: S^d \rightarrow \mathbb{R}$ of degree at most t , the average value of f over S^d equals the average of f on X . Spherical designs have been studied extensively via combinatorics, approximation theory, and other fields.

Let G be an additive abelian group. We say that S (a subset of G) is a *t -independent set* in G if for all non-negative integers k and l with $k+l \leq t$, the sum of k (not necessarily distinct) elements of S does not equal the sum of l (not necessarily distinct) elements of S unless the two sums contain the same terms. This concept extends the well studied concepts of sum-free sets and Sidon sets.

In this talk we give some exact values and asymptotic bounds for the maximum size of a t -independent set in the cyclic group and in other abelian groups. As an application, we show how 3-independent sets can be used to construct spherical 3-designs.

From:

<http://www2.math.binghamton.edu/> - Department of Mathematics and Statistics,
Binghamton University

Permanent link:

<http://www2.math.binghamton.edu/p/seminars/comb/abstract.200203baj>



Last update: 2020/01/29 19:03