2025/09/11 01:13 1/1 Cristina Ruiz (Binghamton)

Cristina Ruiz (Binghamton)

A Stratification of the Middle-level MacPhersonian

Abstract for the Combinatorics Seminar 2005 March 31

The *MacPhersonian* MacP(k,n) is the partially ordered set of all oriented matroids of rank k on the ground set $\{1, 2, ..., n\}$, ordered by $M_1 \ge M_2$ if there is a weak map from M_1 to M_2 . MacP(k,n) can be viewed as a combinatorial analog of the Grassmann manifold G(k,n) of k-planes in \mathbf{R}^n .

The Grassmannian G(k,n) has a type of cell decomposition called a *Schubert cell decomposition*. To define the cells we need to fix some subspaces of \mathbf{R}^n . It is known that for a special Schubert cell decomposition of G(k,n), we can give an explicit combinatorial definition of ``cells for a ``cell decomposition of MacP(k,n). This combinatorial analog of a Schubert cell decomposition of G(k,n) is called a *Schubert stratification* of MacP(k,n).

In studying spectral structures on MacP(k, infinity), I found that another stratification of MacP(n, 2n), based on a different Schubert cell decomposition of G(n, 2n), looks promising. I will show the ideas behind this work in progress.

From:

https://www2.math.binghamton.edu/ - **Department of Mathematics and Statistics, Binghamton University**

Permanent link:

https://www2.math.binghamton.edu/p/seminars/comb/abstract.200503rui

Last update: 2020/01/29 19:03