

# Kurt Luoto

## A matroid-friendly basis for quasisymmetric functions.

### Abstract for the Combinatorics Seminar 2007 September 11

Matroid base polytope decompositions arise in the work of certain algebraic geometers. In 2006, Billera, Jia, and Reiner invented a new invariant  $F(M)$  for matroids in the form of a quasisymmetric function. One motivating application of this invariant is to the study of matroid base polytope decompositions. The mapping of matroids to the algebra of quasisymmetric functions (QSym) behaves as a valuation on matroid base polytopes, and leads to a necessary algebraic condition on their decompositions. Billera, Jia, and Reiner pose a number of questions regarding this relationship. We address some of these questions, obtaining a full characterization for the rank two case.

Along the way, we obtain a novel  $Z$ -basis for QSym that has especially nice properties. For instance this basis has nonnegative integer structure constants and reflects, in addition to the usual grading of QSym by degree, a second grading of QSym that on (the images of) loopless matroids coincides with their matroidal rank.

No familiarity with quasisymmetric functions or matroids is assumed for this talk.

From:

<http://www2.math.binghamton.edu/> - **Department of Mathematics and Statistics,  
Binghamton University**

Permanent link:

<http://www2.math.binghamton.edu/p/seminars/comb/abstract.200709luo>



Last update: **2020/01/29 19:03**