×

## Jae-Ho Lee (Tohoku)

## Nonsymmetric Askey-Wilson Polynomials and Q-Polynomial Distance-Regular Graphs

## Abstract for the Combinatorics Seminar 2015 November 24

Roughly speaking, nonsymmetric Askey–Wilson polynomials are eigenfunctions of the Cherednik-Dunkl operator and form a linear basis of the vector space of the Laurent polynomials in one variable. In this talk, I define a finite sequence of certain Laurent polynomials in one variable, using a Q-polynomial distance-regular graph that contains a Delsarte clique. I prove the orthogonality relations for these polynomials, using a representation for a universal double affine Hecke algebra of rank 1. And I show how my Laurent polynomials can be viewed as a finite, combinatorial analogue of the nonsymmetric Askey–Wilson polynomials.

I will give some basic background concerning Q-polynomial distance-regular graphs and explain the significance of the connection between Q-polynomial distance-regular graphs and the double affine Hecke algebra of rank 1 that I found. This talk does not require any background in Hecke algebras. It will be accessible to any graduate and undergraduate students with a knowledge of linear algebra.

From: https://www2.math.binghamton.edu/ - **Department of Mathematics and Statistics, Binghamton University** 

Permanent link: https://www2.math.binghamton.edu/p/seminars/comb/abstract.201511lee

Last update: 2020/01/29 19:03